	MORTONS NEUROMA				
Typical distribution of pain)	Neuroma in situ	Neuroma removed	1	Normal sized nerve	s Skin closure
WHAT IS A MORTONS NEUROMA	A Morton's neuroma is a soft tissue swelling affecting the small nerves between the toes. Most commonly it affects the 2 <sup>nd</sup> and 3 <sup>rd</sup> and 3 <sup>rd</sup> and 4 <sup>th</sup> toes.				
AIMS OF SURGERY	To remove the affected nerve and swollen soft tissues.				
ANAESTHETIC	Regional anesthesia. Local unaesthetic at the knee or ankle to make the limb numb. This means you can go home the same day.				
OVERVIEW					
Operation time	10-30mins				
Incision placement	On the top of the foot between the affected toes				
Stitches	We use dissolvable stitches where possible.				
When do I go home?	You are usually able to go home the same day. You should have a responsible adult to stay with you.				
Time off work	This will vary according to the type of work you do but typically less than 4 weeks.				
INDICATIONS FOR SURGERY	Pain and limitation of daily activities.				
ALTERNATIVE TREATMENTS	Modification of activity and footwear possibly with a foot support.				
I KEAT MEIN 15	Painkillers				
	Cortisone injections				
GENERAL RISKS OF SURGERY	<ul> <li>Infection (minor</li> <li>Scar problems</li> <li>Postoperative pa</li> <li>Loss of sensation</li> <li>Postoperative sw</li> <li>Failure of the op</li> </ul>	in • • relling • eration	Blood o Anaestl Heart a Chroni	hetic problems uttack / angina c pain / swelling	YOU SHOULD READ THIS LEAFLET IN CONJUNCTION WITH THE PREOPERATIVE INFORMATION BOOKLET
SPECIFIC RISKS RELATED TO THIS OPERATION	<ul><li>Stump neuroma.</li><li>Loss of normal s</li></ul>		the foo		
MORE INFO BY:	Fitter Feet For Life 34 North Street Clapham London SW4 0HD Tel 020 7627 4901 www. Fitterfeet.com				